
## VASSAR COLLEGE | UNDERGRADUATE RESEARCH SUMMER INSTITUTE (URSI) SYMPOSIUM | 2020 MODELING THE VENTILATORY BEHAVIOR OF THE EMBRYOS OF LITTLE SKATES

Connor McShaffrey ('21, Cognitive Science) and John Long (Biology, Cognitive Science, Neuroscience & Behavior)



## **Stages 30-31**

**Pulsate:** Gradual, slow undulation without insertion. **Search & Enter**: Rapid undulation down to tail tip to find and insert into a tendril. **Ventilate:** High frequency undulation within a tendril. **Retract:** An incremental pull-back from tendril relative

to sensitivity, amplitude, and frequency.

**Recoil:** Upon sufficient disturbance, pull the tail back to coil it around the body.

## **Stages 31-32**

**Ventilate (L)**: Ventilate with the body bent laterally at the base of the putative adult caudal fin.


**Return to Tendril**: After a full recoil, return to the same tendril that was being ventilated.

**Freeze:** Upon partial disturbance during ventilation, stop moving and wait before ventilating or moving out of the tendril.


## **Stages 32-33**

**Ventilate Front-Faced**: Ventilate with the body bent sharply at the caudal fin so that the tail tip is looped back over the anterior portion of the body disk. **Select & Enter**: Search & Enter, but with greater efficiency

of selection and insertion.



- Embryos of elasmobranch fishes (skates) and some sharks) develop for up to a year in a leathery egg case, the mermaid's purse.
- An embryo must ventilate by undulating its tail inside one of the four hollow tendrils (or







of the capsule to exchange horns) deoxygenated for oxygenated water.

- The development of the embryo's behavior is intimately tied to changes in its own morphology as well as that of the egg capsule. This interaction has never been systematically investigated.
- We model this dynamic behavioral network as a **finite-state system** (see panel above).
- These are embryos of *Leucoraja erinacea*, a North Atlantic species.



Listen to Connor's narration of the highlights of his research.

Scan this QR code with your phone.

 female lays two egg cases at once, 10 to 35 in one year

each egg case holds one embryo

horns plugged for 1/3 of

gestation one year in capsule

tail pumps inside any horn



De Waal, F. (2016). Are We Smart Enough to Know How Smart Animals Are? WW Norton Co.

Leonard, J.B., Summers, A.P., & Koob, T.J. (1999). Metabolic rate of embryonic little skate, Raja erinacea (Chondrichthyes: Batoidea): the cost of active pumping. Journal of Experimental Zoology, 283(1), 13-18.

Long, J.H., & Koob, T.J. (1997). Ventilating the skate egg capsule: the transitory tail pump of embryonic little skates (Raja erinacea). Bulletin of the Mount Desert Island Biological Laboratory, 36, 117-119.

Maxwell, E.E., Fröbisch, N.B., & Heppleston, A.C. (2008). Variability and conservation in late chondrichthyan development: ontogeny of the winter skate (Leucoraja ocellata). The Anatomical Record, 291(9), 1079-1087.



